
The South Dakota QuitLine 101





NOTE: The term "tobacco" refers to commercially produced tobacco products only and never the traditional tobacco of our Northern Plains American Indians



Tobacco use is the SINGLE most PREVENTABLE cause of death in the U.S.




MORE THAN:

HIV/AIDS
Alcohol
Car accidents
Murders
Suicides

COMBINED!!

Damages Caused by Smoking



Source: USDHHS 2004, 2006, 2012. Note: The condition in red is a new disease that has been causally linked to smoking in this report.

Immune function

Overall diminished health




Priority Populations

- American Indian
- Medicaid Clients
- Pregnant Women
- Spit Tobacco Users
- Youth and Young Adults (18< younger)</p>
- Mental Health & Substance Abuse **

** Newly added population as of January 1, 2015

TOBACCO DEPENDENCE: A 2-PART PROBLEM



Dependence is a two part problem and should address both behavioral **AND** physiological aspects

Fiore MC, Bailey WC, Cohen SJ, et al. (2000). *Treating Tobacco Use and Dependence*. *Clinical Practice Guideline*. Rockville, MD: U.S. Department of Health and Human Services, Public Health Service.





1. Free & Easy Enrollment

2. Free Medications

3. Free Coaching

4. Successful & Evidence Based

How to Enroll



- © Call 1-866-SD Quits
- www.sdquitline.com "Click-To-Call"
- Fax Referral
- © EHR Referral



Referrals and Quit Rate

	Number of	Response Rate		95%
	Number of	to Follow-up		Confidence
	Enrollees	Survey	Quit Rate	Interval
Direct				
Healthcare	501	48.7%	48.8%	42.4% – 55.2%
Provider	301	40.770	40.070	42.4% - 33.2%
Referral				
Passive				
Healthcare	12 072	46.5%	45.5%	44.3% – 46.7%
Provider	13,872	40.5%	45.5%	44.5% – 46.7%
Referral				
Other Referral Source	20,493	45.5%	46.1%	45.1% – 47.1%

Kerkvliet, J. (2014) South Dakota Tobacco QuitLine: Cessation Outcomes of Tobacco Users Referred by Healthcare Providers- 2014. [Research Report] Brookings: South Dakota State University, College of Nursing.

SDQuitLine.com



SERVICES

FAQ

RESOURCES & TIPS

PROVIDERS

SHARE C



Welcome to The South Dakota QuitLine.

If you are ready to quit, thinking about quitting, or know someone who wants to quit... we're ready to help. The South Dakota Quitline offers services designed to help you kick the habit for good. This website has been designed as a resource to help you guit and stay quit. All you have to do is call the QuitLine, or you can "click" and we'll call you! It's that easy and it's free!

There are two ways to enroll:



You Call Us

- Open Monday Friday Zam 11pm Saturday 8am - 5pm CST
- free personal live buit coach
- Free cessation medication
- Enrollment takes approximately 10 15 minutes
- Visit the Services tab for more information.

CALL TO ENROLL NOW!

1.866.SD-QUITS 1.866.737.8487



We Call You

- Click here to submit your information to a Quittine Coach.
- Tell us the best time to call you
- Visit the Services tab for more information.

CLICK TO ENROLL NOW!

REQUEST A CALL Will call you.





Enrollment Call



SOUTH DAKOTA
QuitLine

1.866.SD-QUITS www.SDQuitLine.com

- Gather basic information
- Assess readiness to quit in next 30 days
- Oetermine Eligibility
 - 1. SD Resident
 - 2. Tobacco User
 - Re-enrolling
- 10-15 Min
- Open 7am 11 pm M-F & 8am -5 pm Sat. Voicemail available 24 hrs

How Often Can Someone Re-enroll?

- ©Research indicates that cessation often takes multiple attempts for success.
- ©Re-enrollment is unlimited in the number of times a participant can use the QuitLine, but how often is limited.
- Priority Populations can use the QuitLine every 3 months.

QuitLine Medications

QL participants and their QL Coach may choose *one* of the following for FREE:

- Oup to 8 weeks of Rx Medication (need physician to fax Rx):
 - Zyban (Bupropion)
 - Chantix (Varenicline)
- Up to 8 weeks of Over-the-Counter Nicotine Replacement Therapy:
 - **Patch:** 7 mg, 14 mg, 21 mg
 - **Gum**: 2 mg, 4 mg
 - **Lozenge:** 2 mg, 4 mg

How to Receive Medications

Must enroll and continue in QuitLine phone coaching.

- Participant will need to have their healthcare provider prescribe Zyban or Chantix or NRT if under 18 or pregnant
- © Receive (2) 4-weeks supply of RX medication
- © Receive (4) 2-weeks supply NRT (patch, gum, or lozenge)
- All medications sent through a central pharmacy
- Medication is sent to participant's home address



Dual Therapy for 3+ QL Users

- For those who have used the QuitLine 3+ times and still having trouble quitting!
- Participants need the consent and/or prescription from their physician.
- Optional; not required to use dual therapy

Option 1

NRT Patch + Zyban (8 weeks)

Option 2

NRT Patch + (Gum or Lozenge) (8 weeks)





1.866.SD-QUITS 1.866.737.8487 www.SDQuitLine.com

Continue Coaching

- Typically 5 coaching sessions centered around a Quit Date.
- Sessions scheduled by the participant at their convenience.
- QuitLine Coach calls the participant at scheduled time.

Coaching Sessions 1 & 2 Preparation (Prior to Quit Date)



- Develop a Personalized Quit Plan
- Plan a Quit Date
- Discuss Medication Information
- Oiscuss
 - Triggers
 - © Coping Mechanisms
 - Withdrawal Symptoms





Coaching Session 3 On Quit Date

- Coach Provides Support and Encouragement
- Provides Additional Information on Medication

Coaching Session 4 After Quit Date

- Follow-Up
- Review Quit Strategies
- Address Slips




Approximate Length of Coaching Sessions

Call	Time	
Initial Call (1st call to		
QuitLine by a participant to	10-15 minutes	
enroll in program)		
1 st coaching session	25 minutes	
2 nd - 4 th coaching sessions	15-20 minutes	
5 th coaching session	10-15 minutes	

Coaching Session 5 Graduate!

- Issue GRADUATION LETTER
 - ©Can be used to receive lower health insurance premium!
- © Confirm contact information for follow-up
- 7-month follow-up we have a success rate!!!








1.866.SD-QUITS 1.866.737.8487 www.SDQuitLine.com


QuitLine Coach Requirements

- Minimum of Bachelors degree
- Trained in Counseling and behavior modification
- Mave Motivational Interviewing skills
- Sensitive to cultural diversity (Native Americans)
- Mowledgeable in:
 - Health Issues surrounding tobacco use
 - Nicotine Withdrawal
 - ©Cessation medications

Type of Service



Quit Rates by Cessation Product Type (30 dpp), 2012*



*More people quit on varenicline and bupropion than on NRT - X^2 (3, N = 2433) = 15.548, p = 0.01. Confidence intervals for respective quit rates were: NRT = 36.5% to 43.0%, bupropion=43.7% to 60.3%, varenicline = 43.4% to 49.4%, medication plus NRT = 18.5% to 34.8%, and no cessation product = 34.8% to 48.4%.

Success Rates

Other Websites

- BefreeSD.com
- RethinkTobacco.com
- FindyourPowerSD.org
- facebook.com/SDQuitLine

facebook.com/TobaccoRethinkIt

Goodandhealthy.org



Kayla Magee, RN

Statewide Tobacco Cessation Coordinator

Black Hills Special Services Coop 221 S. Central Ave., Suite 33 Pierre, SD 57501

Phone: (605) 224-6287 ext. 238 Fax: (605) 224-8320

Email: kmagee@tie.net